



## ***C. elegans* SIR-2.1 translocation is linked to a proapoptotic pathway parallel to *cep-1/p53* during DNA damage-induced apoptosis**

Sebastian Greiss, Julie Hall, Shawn Ahmed, et al.

*Genes Dev.* 2008 22: 2831-2842

Access the most recent version at doi:[10.1101/gad.482608](https://doi.org/10.1101/gad.482608)

---

**Supplemental  
Material**

<http://genesdev.cshlp.org/content/suppl/2008/10/17/22.20.2831.DC1.html>

**References**

This article cites 63 articles, 26 of which can be accessed free at:  
<http://genesdev.cshlp.org/content/22/20/2831.full.html#ref-list-1>

**Email alerting  
service**

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#)

---

---

To subscribe to *Genes & Development* go to:  
<http://genesdev.cshlp.org/subscriptions>

---

# *C. elegans* SIR-2.1 translocation is linked to a proapoptotic pathway parallel to *cep-1*/p53 during DNA damage-induced apoptosis

Sebastian Greiss,<sup>1</sup> Julie Hall,<sup>2,3</sup> Shawn Ahmed,<sup>2</sup> and Anton Gartner<sup>1,4</sup>

<sup>1</sup>Wellcome Trust Centre for Gene Regulation and Expression, University of Dundee, Dundee DD1 5EH, United Kingdom;

<sup>2</sup>Department of Genetics, Lineberger Cancer Center, and Department of Biology, University of North Carolina, Chapel Hill, North Carolina 27599, USA

*Caenorhabditis elegans* SIR-2.1, a member of the sirtuin family related to *Saccharomyces cerevisiae* Sir2p, has previously been implicated in aging. The mammalian homolog SIRT1 plays important roles in multiple cellular processes including transcriptional repression and stress response. We show that *sir-2.1* is essential for the execution of apoptosis in response to DNA damage, and that *sir-2.1* genetically acts in parallel to the worm p53-like gene *cep-1*. This novel *cep-1*-independent proapoptotic pathway does not require the *daf-16* FOXO transcription factor. Cytological analysis of SIR-2.1 suggests a novel mechanism of apoptosis induction. During apoptosis SIR-2.1 changes its subcellular localization from the nucleus to the cytoplasm and transiently colocalizes with the *C. elegans* Apaf-1 homolog CED-4 at the nuclear periphery. SIR-2.1 translocation is an early event in germ cell apoptosis and is independent of apoptosis execution and *cep-1*, raising the possibility that SIR-2.1 translocation is linked to the induction of DNA damage-induced apoptosis.

[Keywords: *sir-2.1*; SIR2; sirtuin; *C. elegans*; DNA damage response; apoptosis]

Supplemental material is available at <http://www.genesdev.org>.

Received April 11, 2008; revised version accepted August 19, 2008.

Apoptosis during worm development generally requires the transcriptional induction of *egl-1* encoding for a proapoptotic BH3-only domain protein (Conradt and Horvitz 1998, 1999). EGL-1 then interacts with the anti-apoptotic Bcl2 family member CED-9 that is localized at the outer mitochondrial membrane and forms a complex with CED-4, a protein related to mammalian Apaf-1 (Chen et al. 2000). EGL-1 binding to CED-9 leads to the disruption of the CED-9/CED-4 complex. CED-4 is released, accumulates at the nuclear periphery by binding to SUN-1, oligomerizes, and induces the autoactivation of the caspase CED-3, leading to apoptosis (Yang et al. 1998; Horvitz 1999; Chen et al. 2000; Yan et al. 2004, 2005; Fairlie et al. 2006; Tzur et al. 2006). Within somatic tissues apoptosis only occurs during development.

In the proliferative germline of adult worms multiple pathways are able to trigger apoptosis. Interestingly, only

late pachytene stage meiotic germ cells have the potential to undergo apoptosis (Fig. 1D). While physiological germ cell apoptosis occurs independently of exogenous stimuli and is thought to be necessary for maintaining tissue homeostasis (Gumienny et al. 1999), genotoxic stress can also elicit an apoptotic response in germ cells (Gartner et al. 2000). In DNA damage-induced apoptosis, *egl-1* transcription is induced by a pathway that includes ATL-1, a worm ATR-like PI-3 type protein kinase (Garcia-Muse and Boulton 2005); *Caenorhabditis elegans* MRT-2 and HUS-1, which are part of the so-called 9–1–1 DNA sliding clamp complex (Hofmann et al. 2002); and CLK-2, which acts in a pathway parallel to the 9–1–1 complex (Gartner et al. 2000; Ahmed et al. 2001). While upstream sensors and transducers affect all DNA damage responses including DNA repair, cell cycle arrest, and apoptosis, downstream effectors like *cep-1*, which encodes a primordial worm p53-like protein, are only needed for a subset of responses. *cep-1* is required for ionizing radiation (IR)-induced apoptosis, and for cell cycle arrest and apoptosis in response to UV radiation (Fig. 2D; Derry et al. 2001, 2007; Schumacher et al. 2001; Stergiou et al. 2007).

Interestingly, as in mammals where Apaf-1 and Bcl-2 do not directly interact, the regulation of CED-4 seems

<sup>3</sup>Present address: Laboratory of Molecular Toxicology, National Institute of Environmental Health Sciences, Research Triangle Park, Durham, NC 27709, USA.


<sup>4</sup>Corresponding author.

E-MAIL [a.gartner@dundee.ac.uk](mailto:a.gartner@dundee.ac.uk); FAX 44-1382-385695.

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.482608>.

Greiss et al.

**Figure 1.** *sir-2.1* is required for DNA damage-induced germ cell apoptosis. (A) Worms were irradiated with the indicated doses of ionizing radiation at the late L4 larval stage, and apoptotic corpses were scored by DIC optics after 12, 24, and 36 h. (B) Worms were treated as in A and corpses were scored 24 h after irradiation. (C) Representative pictures of germlines of worms scored in B. Arrowheads indicate apoptotic corpses. (D) Schematic drawing of a *C. elegans* germline arm. Germ cell proliferation occurs in the mitotic zone and cells progress through meiosis as they are moved along the germline. In response to DNA damage cell cycle arrest occurs only in the mitotic zone, while only late pachytene cells are able to undergo apoptosis.



to be more complex than previously thought (Meier and Vousden 2007), especially in germ cell apoptosis (see below). Indeed, it has been shown that CED-4 translocates to the nuclear periphery in irradiated mitotic germ cells without concomitantly inducing germ cell apoptosis (Zermati et al. 2007). Furthermore, we show here that CED-4 is predominantly localized at the nuclear periphery in healthy nonapoptotic late pachytene cells. These results, together with the notion that physiological germ cell death occurs independently of *egl-1* (Gumienny et al. 1999), raise the possibility that additional factors are needed to transmit a proapoptotic signal from mitochondrial CED-9 to perinuclear CED-4 to trigger apoptosis.


*C. elegans* SIR-2.1 is a member of the Sirtuins, a ubiquitous family of NAD<sup>+</sup>-dependent protein deacetylases with members present in virtually every species from archaea to mammals (Brachmann et al. 1995). The founding member, budding yeast Sir2p, is involved in the transcriptional silencing of the mating-type loci (Rine and Herskowitz 1987), telomeres (Gottschling et al. 1990), and rDNA repeats (Bryk et al. 1997; Smith and Boeke 1997). Apart from its role in epigenetic regulation of gene expression, Sir2p has been implicated in the nonhomologous end-joining DNA repair pathway (Martin et al. 1999; Hegde and Klein 2000), as well as in double-strand break repair by homologous recombination (Tamburini and Tyler 2005).

In mammals, seven sirtuins exist, the most widely studied of which, SIRT1, is most closely related to budding yeast Sir2p and *C. elegans* SIR-2.1. Interestingly, while most SIR2 mutant phenotypes in budding yeast are related to transcriptional silencing, mammalian SIRT1 interacts with and deacetylates many nonhistone substrates. In tissue culture-based experiments SIRT1 has been reported to have an anti-apoptotic role in response to genotoxic stress. SIRT1 can deacetylate the p53 tumor suppressor protein, which leads to its inactivation and destabilization (Luo et al. 2001; Vaziri et al. 2001; Langley et al. 2002; Cheng et al. 2003). SIRT1 binds and deacetylates FOXO transcription factors, resulting in differential target gene expression. FOXO3 deacetyla-

tion by SIRT1 leads to the transcriptional repression of proapoptotic Bim, but also to the up-regulation of the stress resistance gene GADD45 (Brunet et al. 2004). SIRT1-mediated deacetylation of FOXO1 represses its proapoptotic activity in prostate cancer cells (Yang et al. 2005). In addition, SIRT1 has been reported to deacetylate Ku70, which in the deacetylated form sequesters the proapoptotic factor Bax from mitochondria, thereby inhibiting apoptosis (Cohen et al. 2004). However, recent studies in knockout mice revealed that the *in vivo* role of SIRT1 in stress-induced apoptosis is less clear (Kamel et al. 2006).

In *C. elegans*, *sir-2.1* overexpression is reported to significantly extend post-mitotic life span, whereas deleting *sir-2.1* results in a modestly shortened life span (Tissenbaum and Guarente 2001; Berdichevsky et al. 2006; Wang and Tissenbaum 2006). Extended longevity mediated by *sir-2.1* overexpression requires DAF-16 a forkhead family transcription factor related to the mammalian FOXOs. DAF-16 is negatively regulated by the conserved *C. elegans* insulin/IGF-1 pathway, and is repressed by the DAF-2 insulin receptor (Tissenbaum and Guarente 2001). The life span extension by *sir-2.1* overexpression is mediated by the *C. elegans* PAR-5 and FTT-2 14-3-3 proteins that interact with both DAF-16 and SIR-2.1 (Berdichevsky and Guarente 2006; Wang et al. 2006). *sir-2.1* is also required for life span extension of *eat-2* worms that are long lived, likely due to caloric restriction (Wang and Tissenbaum 2006). In addition, *sir-2.1* is required for germline silencing of multicopy transgenic arrays (Jedrusik and Schulze 2003) and seems to influence the subcellular localization of the linker histone HIS-24 (H1.1) in the germline (Jedrusik and Schulze 2007).

As part of an ongoing program to screen for novel genes involved in DNA repair and/or damage response signaling we screened through *C. elegans* homologs of genes reported to be involved in these processes and included *sir-2.1*. Here we examine the role of *C. elegans* SIR-2.1 in germline DNA damage response pathways. We show that *sir-2.1* specifically affects DNA damage-



**Figure 2.** *sir-2.1* does not affect the DNA damage response pathway upstream of *cep-1*. (A) Cell cycle arrest following DNA damage is not affected by a *sir-2.1* deletion. The number of nuclei is shown in the bottom right corner of each picture. (B,C) *sir-2.1* is not required for *cep-1*-dependent transcriptional induction of *egl-1* and *ced-13*. Worms were irradiated with 120 Gy 24 h after the L4 stage. RNA was extracted 60, 120, and 240 min after irradiation and *egl-1* and *ced-13* transcript levels were assayed by qRT-PCR as described previously (Schumacher et al. 2005a,b). (D) Genetic pathway connecting DNA damage to apoptosis and cell cycle arrest/DNA repair. (E) *sir-2.1* loss of function does not increase radiation sensitivity. Eggs laid per hour (e/h) and the percentage of surviving embryos (hatch rate, h.r.) with and without IR treatment are indicated.

induced apoptosis in parallel to or downstream from *cep-1*, while not impinging on developmental apoptosis or physiological germ cell apoptosis. Finally, we show that SIR-2.1 is located in germ cell nuclei, but is lost from nuclei undergoing apoptosis. SIR-2.1 nuclear loss is an early event in germ cell apoptosis and occurs independent of apoptosis execution. During apoptosis, SIR-2.1 colocalizes with CED-4, suggesting a functional connection between these proteins.

## Results

### *sir-2.1* is required for DNA damage-induced apoptosis

To address whether *sir-2.1* has a role in DNA damage-induced apoptosis, we  $\gamma$ -irradiated late L4 worms and scored apoptosis 12, 24, and 36 h post-treatment. Apo-

ptosis was nearly completely abolished in *sir-2.1(ok434)* mutants (Fig. 1 A,C). The degree of reduction of apoptosis was close to that observed in *egl-1* or *cep-1* mutants (Fig. 1A). To determine if the apoptotic response might just be delayed, rather than absent, we also assayed worms 48 h after irradiation. At this time point we observed a significantly higher number of corpses in the germlines of *egl-1(n3082)* animals as compared with *sir-2.1(ok434)*, while germlines of wild-type worms essentially disintegrated due to excessive apoptosis under those conditions, confirming that SIR-2.1 is indeed necessary for DNA damage-induced apoptosis (Supplemental Fig. 1). To further corroborate these results, we also introduced DNA double-strand breaks by X-ray treatment confirming the absence of apoptosis under those conditions (Supplemental Fig. 2B). We next wanted to verify that the defect in DNA damage-induced apoptosis is due to *sir-2.1(ok434)* rather than due to a secondary, possibly genetically linked mutation. As to this we tested for DNA damage-induced apoptosis with a second *sir-2.1* loss of function allele *sir-2.1(pk1640::Tc1)* that disrupts the *sir-2.1* catalytic domain and confirmed that DNA damage-induced apoptosis is dramatically reduced (Fig. 1A). SIR-2.1 protein was absent in both mutant alleles of *sir-2.1* (Supplemental Fig. 3). To further confirm a role of *sir-2.1* in DNA damage-induced apoptosis we rescued the *sir-2.1(ok434)* deletion with a construct containing the entire *sir-2.1* expressing operon with GFP fused to the C terminus of *sir-2.1*. We found only one line that rescued the *sir-2.1* defect (Fig. 1B), but this became silenced within three to four generations. While the level of DNA damage-induced apoptosis when averaged was comparable with wild type in the rescue line, we noted higher than normal fluctuations in DNA damage-induced apoptosis when scoring individual germlines, with only ~30% of worms showing evidence of DNA damage-induced apoptosis, which then appeared to occur at higher than wild-type levels (data not shown). Thus, these results suggest that overexpressing *sir-2.1* might lead to enhanced apoptosis, which could explain the difficulty of establishing a stable line.

Given that life span extension by *sir-2.1* overexpression is mediated by *daf-16* (Tissenbaum and Guarente 2001), we wanted to see whether the function of SIR-2.1 in DNA damage-induced apoptosis might also depend on DAF-16. It has previously been reported that DAF-16 may be required for DNA damage-induced apoptosis (Pinkston et al. 2006). We therefore assayed the apoptotic response to DNA damage in three *daf-16* mutants: *daf-16(mu27)*, *daf-16(mu86)*, and *daf-16(mgDf50)*. We found only a slight reduction in DNA damage-induced apoptosis in *daf-16(mu86)* and *daf-16(mgDf50)* at early time points and at intermediate doses of irradiation throughout the time course experiment, whereas apoptosis appeared close to wild-type levels in the latest time points (Supplemental Fig. 4). These results indicate that, unlike SIR-2.1, DAF-16 is not an integral part of the pathway leading to DNA damage-induced apoptosis. Our data differ from those reported by Pinkston et al. (2006), but are in line with results reported by Quevedo et al. (2007). In

Greiss et al.

summary, our results suggest that *sir-2.1* mutants are defective in DNA damage-induced apoptosis, and that the proapoptotic function of *sir-2.1* is independent of *daf-16*.

*sir-2.1* does not overtly affect DNA repair and acts in a pathway parallel to or downstream from *cep-1* to effect DNA damage-induced germ cell apoptosis


Some *C. elegans* mutants that are defective for DNA damage-induced apoptosis also play roles in orchestrating DNA repair and *Sarcomyces cerevisiae* SIR2 mutants are compromised in the nonhomologous end-joining DNA double-strand break repair pathway (Martin et al. 1999; Hegde and Klein 2000). To ask if *sir-2.1* is required for double-strand break repair in *C. elegans* we tested whether repair by homologous recombination or DNA end joining is defective in *sir-2.1*. We first examined the potential role of *sir-2.1* in nonhomologous end joining, a DNA repair pathway that plays a major role in late-stage *C. elegans* embryos, where most somatic cells are arrested in the G1 phase of the cell cycle (Clejan et al. 2006). We confirmed that upon ionizing irradiation of late-stage embryos, progression to the L4 developmental stage is dramatically retarded in end-joining defective *cku-80* mutants. In contrast, we could not find such an effect in wild-type or *sir-2.1(ok434)* mutant worms (Supplemental Fig. 2A). Thus, *sir-2.1* is dispensable for nonhomologous end-joining in *C. elegans*. We also found that *sir-2.1* has no overt role in repair by homologous recombination, which is the predominant form of repair in the germline (Clejan et al. 2006). In contrast to the known checkpoint mutants *mrt-2(e2663)* and *hus-1(op241)* (Ahmed et al. 2001) we found that *sir-2.1* mutants do not have an obvious DNA repair defect upon ionizing irradiation (Fig. 2E). Even though we found that *sir-2.1* worms are not overtly radiation-sensitive, we decided to look more closely at whether *sir-2.1* might act as an upstream checkpoint gene by assessing DNA damage-induced cell cycle arrest, which we found to be indistinguishable from that of wild type (Fig. 2A). Defective DNA damage-induced apoptosis in the presence of IR-dependent cell cycle arrest and DNA repair was also found in *cep-1* mutants (Derry et al. 2001; Schumacher et al. 2001). *cep-1* encodes for the *C. elegans* p53-like transcription factor, which we and others have shown to be required for DNA damage-dependent germ cell apoptosis and the transcriptional induction of the BH3-only domain encoding gene *egl-1* (Derry et al. 2001; Schumacher et al. 2001, 2005a,b). Thus *sir-2.1* might, like *cep-1*, specifically affect DNA damage-induced apoptosis.

Given the phenotypic similarity between *cep-1* and *sir-2.1*, we tested whether *sir-2.1* might be required for *cep-1* activation by comparing the transcriptional level of the *cep-1* targets *egl-1* and *ced-13* in wild-type, *cep-1*, and *sir-2.1* strains after irradiation. These experiments show that *cep-1* activity is not affected by *sir-2.1* (Fig. 2B,C), suggesting that *sir-2.1* acts in a pathway leading to DNA damage-induced germ cell apoptosis in parallel to or downstream from *cep-1*. Alternatively, *sir-2.1* might af-

fect the core apoptosis pathway or only affect germ cell apoptosis without being specific to DNA damage-induced apoptosis.

To assess whether *sir-2.1* affects the core apoptosis pathway used during somatic development we scored developmental apoptosis in early L1 larvae by taking advantage of the *ced-1(e1735)* apoptotic corpse engulfment mutant, which allows for visualization of persisting corpses generated during embryonic development in L1 larvae. We found no difference in the number of apoptotic corpses between *ced-1(e1735)* and *ced-1(e1735); sir-2.1(ok434)* animals, indicating that *sir-2.1* does not affect general apoptosis (Fig. 3C). Consistent with this interpretation, we found no extra (undead) cells in the anterior part of the pharynx in late L3 early L4 *sir-2.1(ok434)* animals as would be expected if apoptosis had failed (Fig. 3D).

To exclude the possibility of the core apoptotic pathway being differentially regulated in the germline, or being differentially affected by *sir-2.1* in the germline, we constructed a double mutant with *ced-9(n1653ts)* (Hengartner et al. 1992; Hengartner and Horvitz 1994). This



**Figure 3.** *sir-2.1* specifically affects only DNA damage-induced germ cell apoptosis. (A) Germ cell apoptosis induced by the *ced-9(n1653ts)* temperature-sensitive loss of function mutation is largely unaffected by loss of *sir-2.1*. Worms were grown at 15°C until the late L4 stage and then shifted to 20°C. Apoptosis was scored 24 h later. The slight reduction in *sir-2.1(ok434); ced-9(n1653ts)* worms is likely due to slightly reduced germ cell proliferation of *sir-2.1(ok434)* worms. (B) *sir-2.1(ok434)* has no effect on excessive physiological germ cell apoptosis conferred by the *gla-3(op216)* mutation. Corpses were scored at 20°C 24 after the L4 stage. (C,D) Developmental apoptosis is not affected by *sir-2.1*. (C) Apoptotic corpses that persist until the early L1 stage in *ced-1(e1735)* animals were scored as described (Ellis et al. 1991). (D) Defects in developmental apoptosis were assayed by counting extra cells in the pharynx of old L3/young L4 larvae as described (Ledwich et al. 2000).

allele of *ced-9* encodes a protein that is unable to repress apoptosis at temperatures above 15°C. The increase in apoptosis observed in *ced-9(n1653ts)* worms was not suppressed by *sir-2.1(ok434)* in comparison with apoptosis suppression upon IR (Fig. 3A). These results indicate that *sir-2.1* does not largely affect the core apoptotic pathway in the germline. The slight apoptosis reduction in *ced-9(n1653ts); sir-2.1(ok434)* worms is likely due to the weak reduction in germ cell proliferation of *sir-2.1(ok434)* worms (Fig. 2E). We next considered a potential role for *sir-2.1* in DNA damage-independent germ cell apoptosis. First, we noted that the basal, radiation, and *cep-1*-independent level of germ cell apoptosis referred to as physiological germ cell apoptosis was not obviously diminished in *sir-2.1* mutants (Fig. 1A). We next asked whether *sir-2.1* affects apoptosis in the *gla-3(op216)* mutant background, which shows increased physiological germ cell apoptosis (Kritikou et al. 2006). Figure 3D indicates that there is no overall reduction of apoptosis in a *gla-3(op216); sir-2.1(ok434)* background, indicating that physiological, *egl-1*-independent apoptosis is unlikely to be affected by *sir-2.1* (Fig. 3B). Taken together, our results suggest that the proapoptotic activity of *sir-2.1* is confined to apoptosis occurring in response to DNA damage.

#### *SIR-2.1 is expressed in germline nuclei and is lost in dying germ cells*

To investigate the localization of SIR-2.1 we expressed recombinant full-length SIR-2.1 protein and generated rabbit and goat anti-SIR-2.1 antibodies (Supplemental Fig. 5). Using these antibodies, we found that SIR-2.1 was present in the nuclei of almost all germ cells (Fig. 4; data not shown). This is consistent with previous reports suggesting that SIR-2.1 is nuclear in nongerm cells (Berdichevsky et al. 2006; Wang and Tissenbaum 2006). However, upon irradiation, SIR-2.1 disappeared from the nuclei of many, but not all late-stage pachytene germ cells, while in corresponding nonirradiated samples only very few cells lost SIR-2.1 nuclear staining (Fig. 4 A, arrows for exemplary cells; Supplemental Movie 1 for scanning through the depicted irradiated germline). The above staining, done with goat anti-SIR-2.1, was confirmed using the rabbit anti-SIR-2.1 antibody (Supplemental Fig. 6; Supplemental Movie 6).


Given that only late-stage pachytene cells die by apoptosis, we next asked whether SIR-2.1 disappearance occurs in dying cells or surviving cells. We aimed at using CED-4 localization as a marker for apoptotic cells. Previous studies of developmental apoptosis have shown that during developmental apoptosis CED-4 accumulates at the nuclear periphery in dying somatic cells (Chen et al. 2000). We generated specific rabbit and goat anti-CED-4 antibodies (Supplemental Fig. 5) and using both antibodies (data not shown) found that CED-4 accumulates at the nuclear periphery of dying germ cells (identified by their intense DAPI staining of condensed chromatin) (Fig. 4A, arrows, panels b,e,h,k). Surprisingly, CED-4 also localized around the nuclei of apparently

healthy cells in irradiated germlines or in untreated germlines albeit with lower intensity (Fig. 4A, arrowhead in panels a,b). In irradiated *sir-2.1* and *cep-1* mutants CED-4 did not accumulate around germ cell nuclei as strongly as in irradiated wild-type worms, consistent with the absence of DNA damage-induced apoptosis in these mutants (Supplemental Fig. 7). Thus, although CED-4 is presumably activated through *cep-1*-dependent *egl-1* transcriptional induction in *sir-2.1* mutants, DNA damage-induced perinuclear hyperaccumulation of CED-4 is disrupted, suggesting that execution of apoptosis depends on the SIR-2.1-dependent recruitment or retention of CED-4 near the nucleus.

Since we observed fewer nuclei showing CED-4 hyperaccumulation as compared with the number of corpses under DIC optics and given that SIR-2.1 was lost from many nuclei without CED-4 hyperaccumulation (Fig. 4B, panels c,i), we suspected that CED-4 perinuclear hyperaccumulation might be a late-stage germ cell apoptosis marker, consistent with the condensed chromatin of those cells (Fig. 4A, panels b,h). We therefore decided to employ a further apoptosis marker. In apoptotic cells mitochondria fragment, leading to the loss of the fine-meshed mitochondrial network and the accumulation of condensed morphologically distinct organelles (Jagasia et al. 2005). We confirmed these changes in mitochondrial morphology by staining irradiated germ cells with a cocktail of commercially available monoclonal antibodies against conserved mitochondrial proteins (Fig. 4B; Materials and Methods). The disappearance of the fine-meshed mitochondrial network and the concomitant appearance of condensed punctiform mitochondria correlated with the loss of nuclear SIR-2.1 (Fig. 4B, arrows; Supplemental Movies 2–5 for scanning through the depicted germline) while the fine-meshed mitochondrial network typical for healthy cells was present in cells with nuclear SIR-2.1 (Fig. 4B, arrowheads; Supplemental Fig. 8). Scanning through irradiated germlines indicates that a fine-meshed mitochondrial network persists in the nucleus free rachis located at the center of the germline, and that the absence of the fine-meshed mitochondrial network best correlates with SIR-2.1 nuclear loss (Supplemental Movies 2–5 for scanning through the depicted germline). In the outer rim of the germline, mitochondria appeared as condensed (and not fine-meshed) elongated structures in nonapoptotic cells (Supplemental Movies 2–5). We next assessed if SIR-2.1 nuclear disappearance is due to nuclear degradation. We scanned many apoptotic nuclei and found that some cells without nucleoplasmic SIR-2.1 contained SIR-2.1 halos around the nucleus, indicating that SIR-2.1 is likely to be translocated from the nuclei of dying cells rather than being degraded within the nucleus (Figs. 4B [empty arrowheads, panels c,d], 5 [panels e,f]; Supplemental Fig. 6; Supplemental Movies 4–6). These halos also indicated that the nuclear envelope was still intact in those cells, as SIR-2.1 was excluded from the nucleoplasm.

We next wished to address whether SIR-2.1 nuclear loss is an early or late apoptotic event by comparing SIR-2.1 loss with the loss of nuclear envelope integrity that

Greiss et al.




**Figure 4.** SIR-2.1 translocates from the nucleus to the cytoplasm of dying cells. Worms were irradiated at the late L4 larval stage and germlines were extracted and fixed 24 h after treatment. Arrows indicate nuclei of dying cells that have lost SIR-2.1 but where the nucleus is still intact. Arrowheads indicate surviving nuclei with strong SIR-2.1 staining. Empty arrowheads indicate cells in the late stages of apoptosis where the nucleus has disintegrated. (A) Worms were stained with anti-SIR-2.1 and anti-CED-4 antibodies and with DAPI. Pictures in the third column are an enlargement of the region indicated by the white rectangle in b. (B) Worms were treated as described in A and stained with anti-SIR-2.1, anti-CED-4, DAPI, and with a mix of five monoclonal antibodies recognizing different mitochondrial proteins. Pictures taken are a projection of a single cell layer. The right column is an enlargement of the region indicated by the white rectangle in panel a. (C) Worms were irradiated as in A and B and stained as indicated. MAb414 was used as a nuclear envelope marker (Lee et al. 2000). Bars, 10  $\mu$ m. Germlines were stained with goat anti-SIR-2.1 (126.3) and rabbit anti-CED-4 (9103.1) antibodies.

occurs at a late stage of apoptosis, by using the MAb414 antibody recognizing the nuclear pore complex (Lee et al. 2000). We found that the majority of SIR-2.1-negative nuclei (Fig. 4C, arrows) had an intact nuclear membrane (Fig. 4C, middle panel). Nuclei that lacked both SIR-2.1 and MAb414 staining tended to occur more proximal in the germline, indicating that these nuclei represented cells in the very late stages of apoptosis, a notion confirmed by the condensed appearance of chromatin and accumulation of CED-4. We thus conclude that SIR-2.1 nuclear disappearance is an early apoptotic event.

We next asked whether SIR-2.1 translocation might be an active regulatory event or a mere consequence of early apoptosis progression. If SIR-2.1 export is regulatory and *sir-2.1* acts genetically in parallel to *cep-1*, SIR-2.1 export should also occur in *cep-1* mutants. We therefore examined SIR-2.1 and CED-4 localization as well as chromatin morphology in late stage pachytene cells in a *cep-1* mutant background where the basal level of germ cell

apoptosis, termed physiological germ cell apoptosis, is not affected, whereas irradiation-induced apoptosis is almost completely blocked. Under these conditions many late pachytene cell nuclei had largely reduced levels of SIR-2.1 even in the absence of CED-4 perinuclear accumulation (Fig. 5, arrows) or changes in mitochondrial morphology (data not shown) while SIR-2.1 was lost from only very few nuclei of unirradiated germ cells (Supplemental Fig. 8). Similar to wild-type dying cells SIR-2.1 accumulated in the cytoplasm around some nuclei that had lost SIR-2.1 staining (Fig. 5, arrows, panels e,f). Interestingly, we occasionally also found apoptotic cells with strong CED-4 staining and condensed chromatin, which we consider to be apoptotic (Fig. 5, empty arrowhead, left panel). These cells may be dying by the *cep-1*-independent physiological germ cell apoptosis pathway or due to damage-induced apoptosis triggered by spontaneous recombination failure or residual *cep-1* activity or endogenous DNA damage (Gumienny et al.



**Figure 5.** SIR-2.1 translocation is independent of *cep-1* and *ced-3*. Worms were treated and germlines were stained as described in Figure 4. Arrows indicate nuclei that have lost SIR-2.1 but where the nucleus is still intact. Arrowheads indicate nuclei with strong SIR-2.1 staining. Empty arrowheads indicate cells in the late stages of apoptosis. Germlines were stained with goat anti-SIR-2.1 (126.3) and rabbit anti-CED-4 (9103.1) antibodies.

1999; Gartner et al. 2000). Consistent with the idea that SIR-2.1 translocation might be linked to an early apoptotic event independent of apoptosis execution, we also observed that SIR-2.1 exits from the nucleus of germlines of irradiated *ced-3(n717)* caspase defective worms that rarely show a single apoptotic corpse (Fig. 5). In contrast, SIR-2.1 remains nuclear in unirradiated germlines (Supplemental Fig. 8).

We next tested if the upstream DNA damage checkpoint mutants *atl-1*, *clk-2*, and *mrt-2* affect SIR-2.1 translocation. We therefore treated *atl-1(tm853)*, *clk-2(qm37)*, and *mrt-2(e2663)* worms with 120 Gy of ionizing irradiation. While we see apoptosis induction and SIR-2.1 nuclear loss in the majority of late pachytene cells in wild-type (Fig. 4; Supplemental Fig. 6; Supplemental Movies 1,2,4,5,6) and in *atl-1/+* heterozygotes (Fig. 6), SIR-2.1 is only lost from a small minority of late pachytene nuclei in *clk-2* and *atl-1* checkpoint mutants, while the reduction in *mrt-2* appears as less dramatic (Fig. 6). We think that there are several reasons why SIR-2.1 translocation is not completely blocked in these strains. We recently showed that DNA damage checkpoint mutants do not fully block *egl-1* induction (Greiss et al. 2008), indicating that a residual level of checkpoint signaling still occurs in those mutants. Furthermore, as checkpoint responses are defective in *mrt-2*, *clk-2*, and *atl-1* germlines (Gartner et al. 2000; Ahmed et al. 2001; Garcia-Muse and Boulton 2005), there are more late pachytene cells in these germlines as compared with wild type due to the lack of apoptosis and cell cycle


arrest. In summary, these data suggest that SIR-2.1 translocation largely depends on the DNA damage checkpoint pathway.

The loss of SIR-2.1 from dying cells raises the possibility that there might be a direct link between SIR-2.1 and apoptosis proteins. However, we could not observe a direct interaction between SIR-2.1 and the *C. elegans* cell death proteins CED-9, CED-4, or EGL-1 by coimmunoprecipitation experiments (data not shown). This may be a consequence of only a small minority of worm cells being in the process of dying at any given time. However, close examination of wild-type irradiated early pachytene nuclei revealed perinuclear dots of accumulated SIR-2.1 that often also showed increased CED-4 staining, most apparent when using goat anti-CED-4 and rabbit anti-SIR-2.1 antibodies (Fig. 7, left panel). We next examined if these structures, which may reflect a very transient colocalization between SIR-2.1 and CED-4 before cells apoptose, might accumulate in apoptosis defective *ced-3(n717)* germ cells. SIR-2.1 and CED-4 were indeed more extensively colocalized in *ced-3(n717)* mutants (Fig. 7, right panel). In summary, our cytological data are consistent with a model that SIR-2.1 translocation and a possible functional interaction with CED-4 might be an integral part of DNA damage-induced apoptosis.

## Discussion

Our studies identified a novel function of *sir-2.1* in promoting DNA damage-induced germ cell apoptosis. This

Greiss et al.




**Figure 6.** SIR-2.1 translocation depends on the DNA damage checkpoint pathway. Worms were irradiated as described in Figure 4, and germlines stained with anti-SIR-2.1 antibodies and DAPI. Arrows indicate nuclei that have lost SIR-2.1. Bar, 10  $\mu$ m.

proapoptotic function of *sir-2.1* is restricted to DNA damage-induced apoptosis, as neither developmental apoptosis nor physiological germ apoptosis nor germ cell apoptosis in *gla-3* and *ced-9* loss-of-function mutants is overtly affected by *sir-2.1*. *sir-2.1* seems to be specifically needed for DNA damage-induced apoptosis, as other DNA damage responses such as transient germ cell cycle arrest and DNA repair are not affected in the mutant. DNA damage signaling, which induces *cep-1*-dependent *egl-1* transcription (Schumacher et al. 2005a,b), is not compromised in *sir-2.1* mutants, suggesting that *sir-2.1* genetically acts in parallel to, or downstream from, *cep-1*-dependent transcription to affect DNA damage-induced apoptosis (Fig. 8).


The finding that *sir-2.1* is required for DNA damage-induced apoptosis is surprising in light of in vitro and tissue culture-based studies that indicate that mammalian SIRT1 suppresses stress-induced apoptosis by deacetylating p53 (Luo et al. 2001; Vaziri et al. 2001). The hypothesis that SIRT1 is a negative regulator of p53 was supported by observation of increased apoptosis in thymocytes from irradiated SIRT1 knockout mice (Cheng et al. 2003). However, recent studies indicate that thymocyte apoptosis is not modulated by SIRT1 and that SIRT1 does not affect transcription of p53 targets even though SIRT1 and p53 can physically interact (Kamel et al. 2006). Our results in *C. elegans* may suggest proapoptotic in vivo function(s) in response to DNA damage for mammalian sirtuins.

Our observations that SIR-2.1 functions independently of *cep-1*-induced transcription of the *egl-1* BH3-only gene, and that DNA damage-induced translocation of SIR-2.1 occurs independently of the core apoptotic machinery are supported by previous studies that apoptosis induction can be affected downstream from or in parallel to *egl-1* and or *ced-9*. For example, *pal-1*-dependent tran-

scriptional induction of *ced-3* occurs during the death of the worm tail spike cell, where *egl-1* and *ced-9* only play minor roles (Maurer et al. 2007). Similarly, the *ceh-30* transcription factor, which regulates the sex-specific death of the CEM neurons, acts downstream from or in parallel to *egl-1* and *ced-9* (Peden et al. 2007; Schwartz and Horvitz 2007). In germline apoptosis it has been reported that the *C. elegans* retinoblastoma gene homolog *lin-35* and the E2F like transcription factor components *efl-2* and *dpl-1* are required for IR-induced germ cell apoptosis independent of *egl-1* regulation, likely through



**Figure 7.** SIR-2.1 colocalizes with CED-4 in germ cells after irradiation. The apparent weak intranuclear CED-4 staining is nonspecific. Bar, 5  $\mu$ m. Germlines were stained with rabbit anti-SIR-2.1 (1434.3) and goat anti-CED-4 (10147.1).



**Figure 8.** Model. SIR-2.1 is exported from germ cell nuclei upon DNA damage and impinges on the apoptotic pathway at the genetic level of EGL-1/CED-9/CED-4.

transcriptional regulation downstream from and/or in parallel to *cep-1*, or through transcriptional regulation of *ced-9*, *ced-4*, and *ced-3* (Schertel and Conradt 2007). In contrast to *lin-35* or *efl-2* and *dpl-1* mutations, we did not find any effect on *ced-9*, *ced-4*, and *ced-3* transcription in *sir-2.1* mutants (data not shown).

How could SIR-2.1 impinge on germ cell apoptosis independent of transcriptional regulation? Our data indicate that it is unlikely that *sir-2.1* affects apoptosis through transcriptional regulation. If the loss of SIR-2.1 from the nucleus was needed for the activation of genes promoting germ cell apoptosis, deletion of *sir-2.1* should phenocopy the effect of SIR-2.1 translocation and lead to excessive rather than blocked apoptosis. Our results indicate that at least in germ cells, apoptosis execution might be more complex than previously thought based on studies on developmental apoptosis. In this system, mitochondrial-bound CED-9 is thought to be complexed to CED-4 to keep CED-4 at bay in order to prevent CED-3 caspase activation. Once *egl-1* is transcriptionally induced EGL-1 releases CED-4, which then translocates to a perinuclear location and triggers CED-3 activation, likely through oligomerization (Yang et al. 1998; Horvitz 1999; Chen et al. 2000). In contrast, we found that CED-4 constitutively localizes around the nuclear membrane in late pachytene stage cells. CED-4 perinuclear hyperaccumulation, which in developmental apoptosis is considered an early event of apoptosis activation (Chen et al. 2000), only occurs in few rather late-stage corpses after SIR-2.1 nuclear exit. Thus, it is unlikely that the direct displacement of CED-4 from the mitochondrial-bound CED-9/CED-4 complex occurs solely by direct EGL-1 binding. The worm germline system might thus be more analogous to mammalian Apaf-1 regulation than previously thought, and may lack a direct physical link between mitochondrially located Bcl-2-like CED-9 and the Apaf-1-like CED-4 to regulate apoptosis (Danial and Korsmeyer 2004). We speculate that one of the factors regulating CED-4 during DNA damage-induced germ cell apoptosis might be SIR-2.1, which appears to transiently colocalize with CED-4, once its translocation from the nucleus to the cytoplasm commences. This regulation is likely not a direct consequence of CED-4 deacetylation, as we could not detect a direct interaction between recombinant CED-4 and SIR-2.1, and coimmu-

noprecipitation from irradiated whole worm extracts failed. Nevertheless, aside from our genetic data, the weak perinuclear colocalization of CED-4 and SIR-2.1, observed in wild-type germlines, which is enhanced in apoptosis execution-defective backgrounds, suggests a functional relationship between these proteins. Future studies will address exactly how DNA damage-induced germ cell apoptosis is triggered and if these mechanisms are conserved in mammals.

## Materials and methods

### *C. elegans strains and maintenance*

Worms were maintained at 20°C on NGM agar plates according to standard protocols, unless otherwise indicated. Alleles used were LG I: *cep-1(lg12501)*, *cep-1(gk138)*, *gla-3(op216)*, *ced-1(e1935)*, *daf-16(mu26)*, *daf-16(mu86)*, *daf-16(mgDf50)*; LG III: *mrt-2(e2663)*, *clk-2(qm37)*, *ced-9(n1653)*, *ced-4(n1162)*, *lig-4(ok716)*, *cku-80(ok861)*, *unc-119(ed3)*; LG IV: *sir-2.1(ok434)*, *sir-2.1(pk1640::Tc1)*, *ced-3(n1717)*, *ced-3(n2438)*; LG V: *atf-1(tm853)*, *egl-1(n1084n3082)*. *sir-2.1(pk1640::Tc1)* carries a Tc1 transposon insertion between the nucleotides corresponding to nucleotides 17,144 and 17,145 of cosmid R11A8.

### *C. elegans apoptosis and DNA damage response assays*

DNA damage-induced apoptosis and radiation sensitivity (rad) assays were performed as described (Gartner et al. 2000). For  $\gamma$ -irradiation a Cs137 source, 2.9 Gy/min (IBL 437C, CIS Bio International) and for X-ray treatment a Stabilipan (Siemens) source was used (11.25 Gy/min). For rad assays, late L4 stage worms were  $\gamma$ -irradiated, transferred to fresh plates 24 h later, and removed after 12 h. Defects in NHEJ repair were scored as described (Clejan et al. 2006). Developmental apoptosis was scored as described previously (Ellis et al. 1991). *egl-1* and *ced-13* quantitative RT-PCR (qRT-PCR) was performed using ~1000 age-synchronized worms using *tbg-1* as an internal normalization control as described previously (Schumacher et al. 2005b). The following primers were used: *tbg-1*: GA1760 (5'-AAGATCTATTGTTCTACCAGGC-3') and GA1761 (5'-CTTGAACCTTCTTGCTCCTTGAC-3'); *egl-1*: GA1762 (5'-CCTCAACCTCTTCGGATCTT-3') and GA1763 (5'-TGCTGATCTCAGAGTCATCAA-3'); *ced-13*: GA1764 (5'-GCTCCCTGTTATCACTTCTC-3') and GA1765 (5'-CTGGCATACGTCTGAATCC-3').

The SIR-2.1::GFP fusion plasmid (pGA291) was constructed by cloning the *C. elegans* operon CEOP4372 including the endogenous promoter and 3'UTR into a vector containing the *C. elegans unc-119* gene as a transformation marker (Praitis et al. 2001). The sequence encoding for GFP and containing artificial introns was amplified from Addgene plasmid 1587 and cloned 5' of the *sir-2.1* stop codon, resulting in a C-terminal GFP fusion. Primers used were GA1263 (5'-GCGTGGGCGGCCATGC GAAAGTTAGACACTAGGCG-3'), GA1351 (5'-AGATGCGGCCGAGATACGCATTTCTTCACACAAA-3'), GA1350 (5'-TCAGTGGCGCCGCTGAATCTCATGTTAAAAAATTTCAA-3'), and GA1299 (5'-CGCAGGCCGGCCCTACCAGCCATGATACTCTACGC-3'). Transgenic lines were created by Biolistic bombardment using a PDS-100/He Biolistic Particle Delivery System (Bio-Rad).

### *Protein expression and antibody production*

6xHis-tagged full-length SIR-2.1 (pGA225) and CED-4 (pGA333), and MBP-tagged SIR-2.1(pGA226) and CED-4 (pGA334) were

Greiss et al.

amplified from cDNA derived from adult worms using the same protocol as used for the qRT-PCR (Schumacher et al. 2005b) using primers GA1346 (5'-TTCAGGCCGGCCTGATACGCA TTTCTTCACACAAA-3') and GA1348 (5'-AACGTGGCGC GCCATGTCACGTGATAGTGGCAAC-3') for *sir-2.1*, and GA1911 (5'-TAACGGCGGCCATGCTCTGCGAAATCGAA TGC-3') and GA1912 (5'-ATCAGGCCGGCCACAGCATG CAAAATTTTTGAGG-3') for *ced-4* to introduce *AscI* at the 5' of the start codon and *FseI* at the 3' of the stop codon and cloned into appropriately modified pQE-80L (6xHis) and pMAL-c2 (MBP). Protein expression was done in BL21(DE3) CodonPlus grown at 37°C to an OD<sub>600</sub> = 0.6 before shifting to 20°C and adding IPTG (1 mM). Bacteria were harvested after incubation for 3 h (SIR-2.1) or overnight (CED-4). 6xHis-tagged SIR-2.1 was soluble and purification was carried out with Ni-NTA (Qiagen). 6xHis-tagged CED-4 was recovered from inclusion bodies using BugBuster (Novagen), solubilized with 10 mM Tris, 300 mM NaCl, 10 mM imidazole, 8 M urea, and purified with Ni-NTA (Qiagen). After elution in the presence of 8 M urea the protein was refolded by stepwise dialysis at 4°C in PBS + 4 M urea (overnight), PBS + 2 M urea (2 h), PBS + 1 M urea (2 h), PBS (2 h) ×2, PBS (overnight). Maltose-tagged proteins were purified according to standard protocols on amylose resin (New England Biolabs). For affinity purification, proteins were covalently linked to AffiGel 15 (Bio-Rad). 6xHis-tagged proteins were used to immunize rabbits and goats. Antibodies were then affinity-purified from the final bleeds using MBP-tagged protein.

#### Immunostaining of isolated *C. elegans* germlines

Worms were dissected on poly lysine coated slides in egg buffer (Edgar 1995) supplemented with 0.1% Tween 20 and 0.2 mM levamisol. Germlines were then fixed in 1.8% formaldehyde for 5 min at room temperature followed by freeze cracking by submersion in liquid nitrogen. Post-fixation was done in a 1:1 mixture of methanol:acetone at -20°C, followed by permeabilization with PBS + 1% Triton X-100 (three times for 10 min, room temperature). Blocking was performed by incubating the samples with Image-iT FX signal enhancer (Invitrogen) for 20 min, followed by 15 min of incubation in PBS + 0.1% Tween 20 + 1% BSA (PBSTB). Primary antibodies were diluted in PBSTB and allowed to bind at 4°C overnight in a humid chamber. Samples were washed three times for 10 min in PBS + 0.1% Tween 20 (PBST). Binding of secondary antibodies was performed for 2 h at room temperature with antibodies diluted in PBSTB supplemented with 1 µg/µL DAPI. After washing three times for 10 min in PBST, the samples were mounted in mounting medium (90% glycerol, 20 mM Tris at pH 8.0, 1 mg/mL p-phenylenediamine). Pictures were taken with a Leica LMF Spectris using SoftWorX software (Applied Precision).

The following primary antibodies were used: goat anti-SIR-2.1 (126.3), 100× dilution; goat anti-CED-4 (10147.1), 250× dilution; rabbit anti-SIR-2.1 (1434.3), 250× dilution; and rabbit anti-CED-4 (9103.1), 100× dilution.

Mitochondria were stained with a 250× dilution (final concentration 0.08 µg/mL for each antibody) of a mixture containing equal amounts of mouse monoclonal antibodies (MitoSciences): MS404 (anti-Complex IV subunit I), MS503 (anti-ATP synthase [Complex V] subunit β), MS507 (anti-ATP synthase [Complex V] subunit α), MSP07 (anti-PDH subunit E1 α), and MSA06 (anti-Cytochrome c). Mouse monoclonal antibody MAb414 (Covance Research Products) was used at a 250× dilution (final concentration 4 µg/mL).

The following secondary antibodies were used for detection: Cy3 labeled donkey anti-rabbit (Jackson Immunochemicals); 1000× dilution, 1.4 µg/mL; FITC labeled donkey anti-goat (Jack-

son Immunochemicals); 100× dilution, 15 µg/mL; FITC labeled donkey anti-mouse (Jackson Immunochemicals); 50× dilution, 28 µg/mL; Alexa 488-labeled donkey anti-goat (Molecular Probes); 200× dilution, 10 µg/mL; Alexa 647-labeled donkey anti-mouse (Molecular Probes); 200× dilution, 10 µg/mL; and Alexa 647 labeled donkey anti-goat (Molecular Probes); 200× dilution, 10 µg/mL.

#### Western blots

Worms were lysed in 4× LDS sample buffer (Invitrogen) by boiling for 15 min. After gel electrophoresis the protein was transferred to a nitrocellulose membrane, blocked with PBST supplemented with 5% milk powder, and probed with antibody 1434.3 (1000× dilution) overnight at 4°C. Horseradish peroxidase-conjugated goat anti-rabbit antibodies (Jackson Immunochemicals) were used as secondary antibodies.

#### Acknowledgments

We are grateful to Julian Blow, Arno Mueller, and Bjoern Schumacher for critical comments on the manuscript; Ehsan Pourkarimi, Caroline Fleischer, and Jean-Luc Tison for help with generating antibodies and immunostaining; and Ronald Plasterk and Marcel Tijsterman for *sir-2.1(pk1640::Tc1)*. A.G. and S.G. were funded by AICR, CRUK (CDA award to A.G.), the DFG, and the Max Planck Society (Erich Nigg). S.A. and J.H. were supported by an Ellison Medical Foundation New Scholar in Aging Award

#### References

- Ahmed, S., Alpi, A., Hengartner, M.O., and Gartner, A. 2001. *C. elegans* RAD-5/CLK-2 defines a new DNA damage checkpoint protein. *Curr. Biol.* **11**: 1934–1944.
- Berdichevsky, A. and Guarente, L. 2006. A stress response pathway involving sirtuins, forkheads and 14–3–3 proteins. *Cell Cycle* **5**: 2588–2591.
- Berdichevsky, A., Viswanathan, M., Horvitz, H.R., and Guarente, L. 2006. *C. elegans* SIR-2.1 interacts with 14–3–3 proteins to activate DAF-16 and extend life span. *Cell* **125**: 1165–1177.
- Brachmann, C.B., Sherman, J.M., Devine, S.E., Cameron, E.E., Pillus, L., and Boeke, J.D. 1995. The SIR2 gene family, conserved from bacteria to humans, functions in silencing, cell cycle progression, and chromosome stability. *Genes & Dev.* **9**: 2888–2902.
- Brunet, A., Sweeney, L.B., Sturgill, J.F., Chua, K.F., Greer, P.L., Lin, Y., Tran, H., Ross, S.E., Mostoslavsky, R., Cohen, H.Y., et al. 2004. Stress-dependent regulation of FOXO transcription factors by the SIRT1 deacetylase. *Science* **303**: 2011–2015.
- Bryk, M., Banerjee, M., Murphy, M., Knudsen, K.E., Garfinkel, D.J., and Curcio, M.J. 1997. Transcriptional silencing of Ty1 elements in the RDN1 locus of yeast. *Genes & Dev.* **11**: 255–269.
- Chen, F., Hersh, B.M., Conradt, B., Zhou, Z., Riemer, D., Grunbaum, Y., and Horvitz, H.R. 2000. Translocation of *C. elegans* CED-4 to nuclear membranes during programmed cell death. *Science* **287**: 1485–1489.
- Cheng, H.L., Mostoslavsky, R., Saito, S., Manis, J.P., Gu, Y., Patel, P., Bronson, R., Appella, E., Alt, F.W., and Chua, K.F. 2003. Developmental defects and p53 hyperacetylation in Sir2 homolog (SIRT1)-deficient mice. *Proc. Natl. Acad. Sci.* **100**: 10794–10799.

- Clejan, I., Boerckel, J., and Ahmed, S. 2006. Developmental modulation of nonhomologous end joining in *Caenorhabditis elegans*. *Genetics* **173**: 1301–1317.
- Cohen, H.Y., Miller, C., Bitterman, K.J., Wall, N.R., Hekking, B., Kessler, B., Howitz, K.T., Gorospe, M., de Cabo, R., and Sinclair, D.A. 2004. Calorie restriction promotes mammalian cell survival by inducing the SIRT1 deacetylase. *Science* **305**: 390–392.
- Conradt, B. and Horvitz, H.R. 1998. The *C. elegans* protein EGL-1 is required for programmed cell death and interacts with the Bcl-2-like protein CED-9. *Cell* **93**: 519–529.
- Conradt, B. and Horvitz, H.R. 1999. The TRA-1A sex determination protein of *C. elegans* regulates sexually dimorphic cell deaths by repressing the *egl-1* cell death activator gene. *Cell* **98**: 317–327.
- Daniel, N.N. and Korsmeyer, S.J. 2004. Cell death: Critical control points. *Cell* **116**: 205–219.
- Derry, W.B., Putzke, A.P., and Rothman, J.H. 2001. *Caenorhabditis elegans* p53: Role in apoptosis, meiosis, and stress resistance. *Science* **294**: 591–595.
- Derry, W.B., Bierings, R., van Iersel, M., Satkunendran, T., Reinke, V., and Rothman, J.H. 2007. Regulation of developmental rate and germ cell proliferation in *Caenorhabditis elegans* by the p53 gene network. *Cell Death Differ.* **14**: 662–670.
- Edgar, L.G. 1995. Blastomere culture and analysis. *Methods Cell Biol.* **48**: 303–321.
- Ellis, R.E., Jacobson, D.M., and Horvitz, H.R. 1991. Genes required for the engulfment of cell corpses during programmed cell death in *Caenorhabditis elegans*. *Genetics* **129**: 79–94.
- Fairlie, W.D., Perugini, M.A., Kvensakul, M., Chen, L., Huang, D.C., and Colman, P.M. 2006. CED-4 forms a 2:2 heterotetrameric complex with CED-9 until specifically displaced by EGL-1 or CED-13. *Cell Death Differ.* **13**: 426–434.
- Garcia-Muse, T. and Boulton, S.J. 2005. Distinct modes of ATR activation after replication stress and DNA double-strand breaks in *Caenorhabditis elegans*. *EMBO J.* **24**: 4345–4355.
- Gartner, A., Milstein, S., Ahmed, S., Hodgkin, J., and Hengartner, M.O. 2000. A conserved checkpoint pathway mediates DNA damage-induced apoptosis and cell cycle arrest in *C. elegans*. *Mol. Cell* **5**: 435–443.
- Gottschling, D.E., Aparicio, O.M., Billington, B.L., and Zakian, V.A. 1990. Position effect at *S. cerevisiae* telomeres: Reversible repression of Pol II transcription. *Cell* **63**: 751–762.
- Greiss, S., Schumacher, B., Grandien, K., Rothblatt, J., and Gartner, A. 2008. Transcriptional profiling in *C. elegans* suggests DNA damage dependent apoptosis as an ancient function of the p53 family. *BMC Genomics* **9**: 334.
- Gumienny, T.L., Lambie, E., Hartwig, E., Horvitz, H.R., and Hengartner, M.O. 1999. Genetic control of programmed cell death in the *Caenorhabditis elegans* hermaphrodite germline. *Development* **126**: 1011–1022.
- Hegde, V. and Klein, H. 2000. Requirement for the SRS2 DNA helicase gene in non-homologous end joining in yeast. *Nucleic Acids Res.* **28**: 2779–2783.
- Hengartner, M.O. and Horvitz, H.R. 1994. *C. elegans* cell survival gene *ced-9* encodes a functional homolog of the mammalian proto-oncogene *bcl-2*. *Cell* **76**: 665–676.
- Hengartner, M.O., Ellis, R.E., and Horvitz, H.R. 1992. *Caenorhabditis elegans* gene *ced-9* protects cells from programmed cell death. *Nature* **356**: 494–499.
- Hofmann, E.R., Milstein, S., Boulton, S.J., Ye, M., Hofmann, J.J., Stergiou, L., Gartner, A., Vidal, M., and Hengartner, M.O. 2002. *Caenorhabditis elegans* HUS-1 is a DNA damage checkpoint protein required for genome stability and EGL-1-mediated apoptosis. *Curr. Biol.* **12**: 1908–1918.
- Horvitz, H.R. 1999. Genetic control of programmed cell death in the nematode *Caenorhabditis elegans*. *Cancer Res.* **59**: 1701s–1706s.
- Jagasia, R., Grote, P., Westermann, B., and Conradt, B. 2005. DRP-1-mediated mitochondrial fragmentation during EGL-1-induced cell death in *C. elegans*. *Nature* **433**: 754–760.
- Jedrussik, M.A. and Schulze, E. 2003. Telomeric position effect variegation in *Saccharomyces cerevisiae* by *Caenorhabditis elegans* linker histones suggests a mechanistic connection between germ line and telomeric silencing. *Mol. Cell. Biol.* **23**: 3681–3691.
- Jedrussik, M.A. and Schulze, E. 2007. Linker histone HIS-24 (H1.1) cytoplasmic retention promotes germ line development and influences histone H3 methylation in *Caenorhabditis elegans*. *Mol. Cell. Biol.* **27**: 2229–2239.
- Kamel, C., Abrol, M., Jardine, K., He, X., and McBurney, M.W. 2006. SirT1 fails to affect p53-mediated biological functions. *Aging Cell* **5**: 81–88.
- Kritikou, E.A., Milstein, S., Vidalain, P.O., Lettre, G., Bogan, E., Doukoumetzidis, K., Gray, P., Chappell, T.G., Vidal, M., and Hengartner, M.O. 2006. *C. elegans* GLA-3 is a novel component of the MAP kinase MPK-1 signaling pathway required for germ cell survival. *Genes & Dev.* **20**: 2279–2292.
- Langley, E., Pearson, M., Faretta, M., Bauer, U.M., Frye, R.A., Minucci, S., Pelicci, P.G., and Kouzarides, T. 2002. Human SIR2 deacetylates p53 and antagonizes PML/p53-induced cellular senescence. *EMBO J.* **21**: 2383–2396.
- Ledwich, D., Wu, Y.C., Driscoll, M., and Xue, D. 2000. Analysis of programmed cell death in the nematode *Caenorhabditis elegans*. *Methods Enzymol.* **322**: 76–88.
- Lee, K.K., Gruenbaum, Y., Spann, P., Liu, J., and Wilson, K.L. 2000. *C. elegans* nuclear envelope proteins emerin, MAN1, lamin, and nucleoporins reveal unique timing of nuclear envelope breakdown during mitosis. *Mol. Biol. Cell* **11**: 3089–3099.
- Luo, J., Nikolaev, A.Y., Imai, S., Chen, D., Su, F., Shiloh, A., Guarente, L., and Gu, W. 2001. Negative control of p53 by Sir2 $\alpha$  promotes cell survival under stress. *Cell* **107**: 137–148.
- Martin, S.G., Laroche, T., Suka, N., Grunstein, M., and Gasser, S.M. 1999. Relocalization of telomeric Ku and SIR proteins in response to DNA strand breaks in yeast. *Cell* **97**: 621–633.
- Maurer, C.W., Chiorazzi, M., and Shaham, S. 2007. Timing of the onset of a developmental cell death is controlled by transcriptional induction of the *C. elegans* *ced-3* caspase-encoding gene. *Development* **134**: 1357–1368.
- Meier, P. and Vousden, K.H. 2007. Lucifer's labyrinth—Ten years of path finding in cell death. *Mol. Cell* **28**: 746–754.
- Peden, E., Kimberly, E., Gengyo-Ando, K., Mitani, S., and Xue, D. 2007. Control of sex-specific apoptosis in *C. elegans* by the BarH homeodomain protein CEH-30 and the transcriptional repressor UNC-37/Groucho. *Genes & Dev.* **21**: 3195–3207.
- Pinkston, J.M., Garigan, D., Hansen, M., and Kenyon, C. 2006. Mutations that increase the life span of *C. elegans* inhibit tumor growth. *Science* **313**: 971–975.
- Praitis, V., Casey, E., Collar, D., and Austin, J. 2001. Creation of low-copy integrated transgenic lines in *Caenorhabditis elegans*. *Genetics* **157**: 1217–1226.
- Quevedo, C., Kaplan, D.R., and Derry, W.B. 2007. AKT-1 regulates DNA-damage-induced germline apoptosis in *C. elegans*. *Curr. Biol.* **17**: 286–292.
- Rine, J. and Herskowitz, I. 1987. Four genes responsible for a position effect on expression from HML and HMR in *Saccharomyces cerevisiae*. *Genetics* **116**: 9–22.
- Schertel, C. and Conradt, B. 2007. *C. elegans* orthologs of components of the RB tumor suppressor complex have distinct

Greiss et al.

- pro-apoptotic functions. *Development* **134**: 3691–3701.
- Schumacher, B., Hofmann, K., Boulton, S., and Gartner, A. 2001. The *C. elegans* homolog of the p53 tumor suppressor is required for DNA damage-induced apoptosis. *Curr. Biol.* **11**: 1722–1727.
- Schumacher, B., Hanazawa, M., Lee, M.H., Nayak, S., Volkman, K., Hofmann, E.R., Hengartner, M., Schedl, T., and Gartner, A. 2005a. Translational repression of *C. elegans* p53 by GLD-1 regulates DNA damage-induced apoptosis. *Cell* **120**: 357–368.
- Schumacher, B., Schertel, C., Wittenburg, N., Tuck, S., Mitani, S., Gartner, A., Conradt, B., and Shaham, S. 2005b. *C. elegans ced-13* can promote apoptosis and is induced in response to DNA damage. *Cell Death Differ.* **12**: 153–161.
- Schwartz, H.T. and Horvitz, H.R. 2007. The *C. elegans* protein CEH-30 protects male-specific neurons from apoptosis independently of the Bcl-2 homolog CED-9. *Genes & Dev.* **21**: 3181–3194.
- Smith, J.S. and Boeke, J.D. 1997. An unusual form of transcriptional silencing in yeast ribosomal DNA. *Genes & Dev.* **11**: 241–254.
- Stergiou, L., Doukoumetzidis, K., Sendoel, A., and Hengartner, M.O. 2007. The nucleotide excision repair pathway is required for UV-C-induced apoptosis in *Caenorhabditis elegans*. *Cell Death Differ.* **14**: 1129–1138.
- Tamburini, B.A. and Tyler, J.K. 2005. Localized histone acetylation and deacetylation triggered by the homologous recombination pathway of double-strand DNA repair. *Mol. Cell Biol.* **25**: 4903–4913.
- Tissenbaum, H.A. and Guarente, L. 2001. Increased dosage of a *sir-2* gene extends lifespan in *Caenorhabditis elegans*. *Nature* **410**: 227–230.
- Tzur, Y.B., Margalit, A., Melamed-Book, N., and Gruenbaum, Y. 2006. Matefin/SUN-1 is a nuclear envelope receptor for CED-4 during *Caenorhabditis elegans* apoptosis. *Proc. Natl. Acad. Sci.* **103**: 13397–13402.
- Vaziri, H., Dessain, S.K., Ng, E.E., Imai, S.I., Frye, R.A., Pandita, T.K., Guarente, L., and Weinberg, R.A. 2001. hSIR2(SIRT1) functions as an NAD-dependent p53 deacetylase. *Cell* **107**: 149–159.
- Wang, Y. and Tissenbaum, H.A. 2006. Overlapping and distinct functions for a *Caenorhabditis elegans* SIR2 and DAF-16/FOXO. *Mech. Ageing Dev.* **127**: 48–56.
- Wang, Y., Oh, S.W., Deplancke, B., Luo, J., Walhout, A.J., and Tissenbaum, H.A. 2006. *C. elegans* 14–3–3 proteins regulate life span and interact with SIR-2.1 and DAF-16/FOXO. *Mech. Ageing Dev.* **127**: 741–747.
- Yan, N., Gu, L., Kokel, D., Chai, J., Li, W., Han, A., Chen, L., Xue, D., and Shi, Y. 2004. Structural, biochemical, and functional analyses of CED-9 recognition by the proapoptotic proteins EGL-1 and CED-4. *Mol. Cell* **15**: 999–1006.
- Yan, N., Chai, J., Lee, E.S., Gu, L., Liu, Q., He, J., Wu, J.W., Kokel, D., Li, H., Hao, Q., et al. 2005. Structure of the CED-4–CED-9 complex provides insights into programmed cell death in *Caenorhabditis elegans*. *Nature* **437**: 831–837.
- Yang, X., Chang, H.Y., and Baltimore, D. 1998. Essential role of CED-4 oligomerization in CED-3 activation and apoptosis. *Science* **281**: 1355–1357.
- Yang, Y., Hou, H., Haller, E.M., Nicosia, S.V., and Bai, W. 2005. Suppression of FOXO1 activity by FHL2 through SIRT1-mediated deacetylation. *EMBO J.* **24**: 1021–1032.
- Zermati, Y., Mouhamad, S., Stergiou, L., Besse, B., Galluzzi, L., Boehrer, S., Pauleau, A.L., Rosselli, F., D'Amelio, M., Amendola, R., et al. 2007. Nonapoptotic role for Apaf-1 in the DNA damage checkpoint. *Mol. Cell* **28**: 624–637.